

Bioneer Corporation

8-11 Munpyeongseo-ro, Daedeok-gu, Daejeon, 34302
Republic of Korea
Tel: +82-42-930-8777 (Korea): 1588-9788
Fax: +82-42-930-8688 E-mail: sales@bioneer.com

Bioneer Inc.

1301 Marina Village PKWY, Suite 110
Alameda, CA 94501, USA
Toll Free: +1-877-264-4300 Fax: +1-510-865-0350
E-mail: order.usa@bioneer.cus.com

Bioneer R&D Center

Korea Bio Park BLDG #B-702, 700 Daewangpangyo-ro
Bundang-gu, Seongnam-si, Gyeonggi-do, 13488
Republic of Korea
Tel: +82-31-628-0500 Fax: +82-31-628-0555

[Cat. No.]
D-1020

[Lot No.]

[Volume]
250 μ l

[Concentration]
150 ng/ μ l



Consult
Instruction
For Use



Research
Use
Only



Catalog
Number



Batch
Code



Contains
Sufficient for
(n) tests



Caution,
Consult
Accompanying
documents



USE BY



Temperature
Limitation


Note

For research use only. Not
for use in diagnostic or
therapeutic procedures.

Bioneer shall not be in any
event be liable for
incidental or special
damage of any kind
resulting from any use
except on application(s).

- **Description** : 25/100 bp Mixed DNA Ladder is specially designed for determining the size of double strand DNA from 25 to 2,000 base pairs. The DNA Ladder consists of 17 double strand DNA fragments ranging in size from 25 to 200 bp in 25 bp increments, and additional fragments of 300, 400, 500, 600, 700, 800, 900, 1000, 2000 bp. The 150, 500, 1,000 and 2,000 bp bands are approximately two to three times brighter for easy identification.

- **Application** : Size confirmation of DNA fragments.



• **Features**

- Ready-to-use
- Recommended Loading:
2 μ l / 5 mm lane width
- Typical Loading Number of Lanes:
125 (5 mm lane width)
- Range (bp): 25 ~ 2,000
- Number of Bands: 17

• **Storage Buffer**

- 10 mM Tris-HCl (pH 8.0)
- 1 mM EDTA
- 2.5% Ficoll
- 0.005% Bromphenol Blue
- 0.005% Xylene Cyanol

• **Recommend Electrophoresis Condition**

- 2.5% agarose gel.
- 0.5X TBE buffer.
- Agarose use for small DNA analysis.

• **Storage Condition**

- This product is shipped on dry ice and should be stored at -20°C until use.

• **Quality control assay**

- Agarose gel analysis shows that 10 bands are clearly distinguishable. The 2,961 bp band must be more intense than any other band.

• **Note**

- The DNA ladder can be applied directly onto an agarose gel.
- Do not heat before loading.
- Repeated freezing and thawing should be avoided.
- If TAE buffer is reused several times, DNA bands occasionally appear smeared in agarose gel electrophoresis. Because the buffering capacity of TBE buffer is higher, it can be used longer than TAE buffer.

• **Ordering Information**

Cat. No.	Product
D-1010	10 bp DNA Ladder, 100 μ l (522 ng/ μ l)
D-1020	25/100 bp Mixed DNA Ladder, 250 μ l (150 ng/ μ l)
D-1030	100 bp DNA Ladder, 250 μ l (135 ng/ μ l)
D-1035	100 bp Plus DNA Ladder, 500 μ l (80.7 ng/ μ l)
D-1040	1 kb DNA Ladder, 500 μ l (130 ng/ μ l)
D-1050	Lambda DNA / Hind III Marker, 500 μ l (200 ng/ μ l)
D-1060	Lambda DNA / EcoR I Marker, 500 μ l (200 ng/ μ l)
D-1070	Lambda DNA / EcoR I + Hind III Marker, 500 μ l (200 ng/ μ l)
D-2510	Lambda DNA, 500 μ l (250 μ g)
D-1030-1	AccuLadder 100 bp DNA Size Marker, 320 μ l x 2 (54 ng/ μ l)
D-1040-1	AccuLadder 1 kb DNA Size Marker, 500 μ l x 2 (65 ng/ μ l)